

Control of grape berry moth larvae using parasitoids: should it be developed?

D. Thiéry¹, L. Delbac¹, C. Villemant², J. Moreau³

¹UMR Santé Végétale 1065 INRA, Institut de Sciences de la Vigne et du Vin de Bordeaux, IFR 103. ²Muséum National d'Histoire Naturelle, Département Systématique & Evolution, UMR7205 OSEB, CP50, Paris, France, ³Équipe Écologie Évolutive, UMR CNRS 5561 Biogéosciences, Université de Bourgogne, F-21000 Dijon, France.

Abstract: Besides mating disruption techniques and progress in monitoring techniques (e.g. the use of food traps against females), biological control may reveal itself very efficient at controlling grape moth populations. Parasitoids active to control grape moths are known for long in vineyards; few of them were already described in the middle of the 19th century in French vineyards and their efficiency was already recognized especially against the diapausing and the first spring generations of the moths. Rather numerous attempts to release egg parasitoids have been done in different European countries using different species of trichogrammas. The results obtained varied a lot and could not yet clearly promote the use of this technique in vineyards. We believe that a biological control based on larval parasitoids could efficiently be developed as a valuable alternative to chemical control. In the present paper, we focus on larval parasitoids among which ichneumonids and chalcidoids (Hymenoptera) dominate, and present results obtained in different French vineyards (Bordeaux vineyard, Perpignan and Montpellier area, Côtes du Rhône and Alsace). We discuss factors that may favour or reduce their efficiency as biocontrol agents.

Key words: Parasitoids, biological control, viticulture, biodiversity, fitness, grape cultivars, Ichneumonids, Tachinids, Trichogrammas.

Introduction

Biological control in vineyard is a promising but challenging perspective, and surprisingly the development of biological control in viticulture suffers from a lack of studies. The main efforts have been done studying and attempting egg parasitoids releases (namely *Trichogrammas*) (see Castaneda-Samoya *et al.*, 1993; Reda Abd el Monsef, 2004 and Hommay *et al.*, 2010). Beside such studies, several others attempted to qualify the biodiversity of natural enemies in different European vineyards (Thiéry, 2008). This started in the middle of the 19th century (Audouin, 1842; Jolicoeur, 1894) until more recent years (e.g. Coscolla, 1980; Marchesini & Dalla Monta, 1994; Colombera *et al.*, 2001; Thiéry *et al.*, 2001; Barnay *et al.*, 2001). Also the biology, ecology or behaviour of several parasitoids occurring in vineyards have been studied both in laboratories and field during these last years focussing on *Trichogrammas* (Stengel *et al.*, 1977; Le Rallec & Wajnberg, 1990; Hommay *et al.*, 2002; Moreau *et al.*, 2009) or larval parasitoids (Chuche *et al.*, 2006; Xuéreb & Thiéry, 2006).

Rather numerous attempts to release egg parasitoids have been done in different European countries using different species of *Trichogrammas* which harvested very different efficacies. Currently the release of parasitoids in vineyards is very marginal, and biological control programmes at large scale in vineyards would require research and development attempting to identify the suitable candidate species to be used like the environmental

conditions favouring these species. This would also require fine basic research concerning the physiology, ethology and ecology of the selected candidates.

Although grape training and pesticide practices may strongly affect the natural enemies population dynamics, most of the field studies concluded that parasitism (either egg or larval) varies according to ecological factors, *e.g.* climatic mismatch, variable host density, lack of alternative hosts, also pointing out that variable quality of the host affect the parasitoid reproductive success. In this paper, we present several ecological parameters influencing the parasitism efficacy.

Parasitoids and predators of grape moths in vineyards

Natural enemies of pest in vineyards have received interest for many years. During the middle of the 19th century, several published monographies already listed eggs or larval parasitoids, the main focus being put on the two main moth pests in French vineyards at this period, the grape berry moth (*Eupoecilia ambiguella*) and the leaf rolling tortrix (*Sparganothis pilleriana*). In a recent review, Thiéry (2008) listed more than 70 species mainly represented in Hymenoptera which shows that vineyard is far from being a 'no parasitoid's land'. In fact parasitoids abundance and diversity may vary a lot throughout the season, between the different grape production areas but also as a consequence of large range of different ecological factors (host density, alternative hosts, grass or floral cover, hedges).

Neighbouring natural landscapes probably play an important role in natural enemies population level (Genini, 2000), even though we miss accurate data from vineyards to properly consider their effect. Interestingly, also natural enemies' biodiversity exists in most of the European vineyards. For example, several oophagous parasitoids as trichogrammas, or larval parasitoids as the tachinid *Phytomyia nigrina* and various hymenopteras as *Campoplex capitator*, *Dibrachys* spp (*cavus*, *affinis*), *Dicaelotus inflexus*, *Diadegma fenestralis*, *Itoplectis maculator*, *Scambus elegans* or *Exochus* spp. (*tibialis*, *notatus*), occur and are regularly found in different European vineyards (Table 1).

Several factors affect the parasitism efficacy

Latitude, climate match or mismatch and vineyard location

These factors are probably not enough investigated. However, in a recent elegant study, Moreau *et al.* (2010) sampled *Lobesia botrana* larvae in different vineyards from Alsace, Switzerland and east South France and compared the larval parasitoids occurrences (Table 2). This shows that both diversity and major species vary according to both latitude and climate, though grape cultivars also contribute to these differences (see below).

This work matches previous studies which also suggest that parasitoids species distribution vary in France according to latitude and climate (Thiéry *et al.*, 2001). Several species are characteristic of Mediterranean viticulture, like for example *P. nigrina* or also *C. capitator* which is classically found in most of Mediterranean viticulture countries though this species has a broader distribution also occurring for example in Switzerland. Because vineyards present a great diversity of climatic conditions, climate match or mismatch is of primary importance and conditions the success of certain parasitoids. During these 2 successive years of study, the species *Exochus tibialis* was exclusively found in Switzerland and Alsace, and dominant in 2003 in the Valaisan vineyards (Table 2), while *P. nigrina* occurred exclusively in the south of France (Côtes du Rhône). In the study conducted by Moreau *et al.* (2010), almost 3 times fold less parasitoids were found in 2004, as compared to 2003, but the vineyards sampled were different.

Table 1. Non exhaustive list of parasitoids (alphabetic order) reported from literature as natural enemies of grape moths in vineyards in West European countries. This list is selected and implemented from a broader one (Thiéry, 2008). Only species reported by at least 2 references in the former list are presented here. EA = *Eupoecillia ambiguella*, LB = *Lobesia botrana*, SP = *Sparganothis pilleriana*, Beth = Hym. Bethyridae, Chal = Hym Chalcidoidea; Ichn = Hym. Ichneumonidae, Tach = Dip. Tachinidae. *Pteromalus* and *Trichogramma* spp regroup several species.

Species	Family	Host orders	Reported hosts in vineyardsI	Parasitized instars (when known).
<i>Agrothereutes abbreviatus</i> (F.)	Ichn	Lepidoptera	LB EA	pupae
<i>Ascogaster quadridenata</i> (Wesm.)	Ichn	Tortricidae	LB	larvae pupae
<i>Brachymeria minuta</i> (Wesm.)	Chal	Lep, Dip	EA	
<i>Campoplex capitator</i> (Aub.)	Ichn	Tortricidae	LB EA	L3-L4
<i>Diadegma fenestrata</i> (Holm.)	Ichn	Lepidoptera	LB SP	larvae
<i>Dibrachys affinis</i> (Masi)	Chal	Lepidoptera and other insects	LB-EA-SP	L4-L5
<i>Dibrachys cavus</i> (Walk.) (syn <i>boucheanus</i>)	Chal	Lepidoptera and other insects	LB-EA-SP	L4-L5 pupae
<i>Dicaelotus inflexus</i> (Thom.)	Ichn	Lepidoptera	LB	pupae
<i>Dicaelotus resplendens</i> (Holm.)	Ichn	Lepidoptera		
<i>Elachertus affinis</i> (Masi)	Chal	Tortricidae	EA-LB-SP	larvae
<i>Exochus tibialis</i> (Holm.)	Ichn	Lepidoptera	LB	larvae pupae
<i>Gelis areator</i> (Panz.)	Ichn	Lep, Hym	LB EA	larvae
<i>Goniozus claripennis</i> (Först.)	Beth	Lepidoptera	PS	larvae
<i>Ischnus alternator</i> (Grav.)	Ichn	Lepidoptera	LB	pupae
<i>Itoplectis alternans</i> (Grav.)	Ichn	Lep, Hym	LB, SP	pupae
<i>Itoplectis maculator</i> (Fabr.)	Ichn	Lep, Hym	LB EA SP	pupae
<i>Itoplectis tunetana</i> (Schmied.)	Ichn	Lep, Hym	L EA	pupae
<i>Phaeogenes melanogonos</i> (Gmel.)	Ichn	Lepidoptera	EA SP	pupae
<i>Phaeogenes planifrons</i> (Wesm.)	Ichn	Lepidoptera	SP	pupae
<i>Phytomyza nigrina</i> (Meig.) (= <i>nitidiventris</i>)	Tach	Lepidoptera	LB	larvae
<i>Pimpla spuria</i> (Grav.)	Ichn	Lepidoptera	LB	pupae
<i>Pimpla turionellae</i> (L.)	Ichn	Lepidoptera	LB EA SP	pupae

Species	Family	Host orders	Reported hosts in vineyardsI	Parasitized instars (when known).
<i>Pteromalus</i> spp <i>More than 8 species</i>	Ichn	Lep, Dip, Col, Hym	LB EA SP	larvae pupae
<i>Scambus elegans</i> (Woldst.)	Ichn	Lep, Hym	LB	larvae
<i>Tranosemella praerogator</i> (L.)	Ichn	Lepidoptera	LB EA SP	larvae
<i>Trichogramma</i> spp <i>More than 15 species</i>	Chal	Lepidoptera	EA LB SP	eggs
<i>Triclistus</i> sp .	Chal	Lepidoptera	LB	larvae pupae

Host density

Host density is a classical major factor affecting natural enemy populations. It becomes crucial for example in specialist parasitoids which lack alternative hosts to achieve their reproductive cycle (Lane *et al.*, 1999). The effect of host density on the parasitoids/predators population level and thus capacities to control the targeted pest is thus well documented in many biological antagonists. With most parasitoids searching for specific development stages, this host density should be viewed at the specific targeted development stage. Xuéreb & Thiéry (2006) showed for example a good correlation between the parasitism by natural populations of *C. capitator* and the number of *L. botrana* larvae per bunch (Figure 1).

Also the host searching behaviour of parasitoids is often affected by host density in the patch. For example, the egg discovery rate of *L. botrana* by *T. cacoeciae* varies as a function of host density (Hommay *et al.*, 1999).

Season of the year and variation between years

Except for the monovoltin leaf rolling tortrix, the grape moths (*L. botrana*, *E. ambiguella* and *Argyrotenia pulchellana*) accomplish at least two generations per year, and their population level may vary sometimes a lot from one generation to another. Therefore and because host density (see above) is of primary importance, parasitoids populations may suffer or profit from these variations.

Variation between successive years is also a trend that has been often observed in different vineyards. This can also be illustrated by the data provided by sampling in 2003 and 2004 *L. botrana* larvae from different vineyards in France and Switzerland (Moreau *et al.*, 2010) (Table 2), but also by data from Xuéreb & Thiéry (2006) (Table 3).

Variation of parasitoids/predators population density may also be due only to their intrinsic annual life cycle or to their life style. Such within year variation is known for long in *C. capitator*. It is worth noting that this species named one century ago *C. majalis* meaning *Campoplex* of may (Audouin, 1842; Jolicoeur, 1894) can diapause in diapausing berry moths chrysalids (*L. botrana* and *E. ambiguella*)(our unpublished data). This species has traditional higher efficiency during the first generation of the moth. These observations were confirmed by Xuéreb & Thiéry (2006) (Table 3). In the same study, *S. elegans* was only found during 2 successive years during summer and never parasitized the spring generation of *L. botrana*.

Intrinsic variation due to the parasitoid life cycle combined with variation of host population level within the year may also amplify such variations.




Figure 1: Host density relationship with the number of *Campoplex capitator* parasitizing the *Lobesia botrana* larvae (from Xuéreb & Thiéry, 2006, see this reference for more details).

Table 2. Species diversity in larval parasitoids emerged from *Lobesia botrana* sampled in different French and Swiss vineyards. VS = Valais, VD = Vaud, F = France. Parasitoid species, Ichneumonidae: *Et* = *Exochus notatus*, *Aa* = *Agrypon anxium*, *Cc* = *Campoplex capitator*, *Im* = *Itopectis maculator*, *Di* = *Dicaelotus inflexus*, *Df* = *Diadegma fenestratale*, Braconidae: *As* = *Apanteles* sp., Bethyliidae: *Gc* = *Goniozus claripennis*, Tachinidae: *Pn* = *Phytomyza nigra*, *Triclistus meridiator* was found only once

Year	Locality	Cultivar	<i>Et</i>	<i>Aa</i>	<i>Cc</i>	<i>Im</i>	<i>Di</i>	<i>Pn</i>	<i>Gc</i>	<i>Df</i>	<i>As</i>
2003	Yvorne (VD)	Pinot noir	6	1	2						
	Yvorne (VD)	Chasselas	7		3						
	St P-de Clages (VS)	Pinot noir	18								
	St P-de Clages (VS)	Gamay	8	1							
	St P-de Clages (VS)	Chasselas	20	2	2						
	Tavel (F)	Grenache			6		1				
	Colmar (F)	Gewurtz	3			2	1				
	Colmar (F)	Riesling				1	3				
Total 2003			62	4	13	3	5	0	0	0	0
2004	Roquemaure (F)	Grenache						16			1
	Roquemaure (F)	Syrah		1	1				1	5	
	Sion (VS)	Pinot noir									
	Sion (VS)	Chasselas	2								
	Nyon (VD)	Chasselas									
	Nyon (VD)	Chardonnay	3								1
Total 2004			5	1	1	0	0	16	1	5	2

Effect of grape cultivars and larval food plant

Grape cultivar, but also host plant consumed as a larva by *L. botrana* which is a polyphagous species (Thiéry & Moreau, 2005; Maher & Thiéry, 2006) surprisingly influenced the susceptibility of its eggs to be parasitized by *Trichogramma evanescens* (Moreau *et al.*, 2009; and Figure 2). Interestingly this study also showed that the reproductive success of the

Trichogrammas emerging from the different eggs also varied like its larval growth speed. A similar result was found with eggs produced by moth fed as larvae on food supplemented with different host-plant of *L. botrana* and exposed to *Trichogramma cacoeciae* (Thiéry, Pizzol & Wanjberg, unpublished data).

Table 3. Parasitism rates by natural populations of *Campoplex capitator* against *Lobesia botrana* during the first 2 successive generations of the year. See Xuéreb & Thiéry, 2006 for details on the procedure.

Parasitoid/cultivar	2000		2001		Mean Pr per cultivar
	G1	G2	G1	G2	
All parasitoids					
Cabernet Franc	50.00	19.05	100.00	21.01	47.51
Cabernet Sauvignon	66.67	34.62	30.00	24.56	38.96
Merlot	20.00	30.67	25.00	31.16	26.71
Sauvignon	36.36	26.67	33.33	20.28	29.16
Sémillon	33.33	33.33	77.78	17.20	40.41
Mean Pr per collection	41.27	28.87	53.22	22.84	
<i>Campoplex capitator</i>					
Cabernet Franc	50.00	4.76	100.00	19.33	43.52
Cabernet Sauvignon	66.67	11.54	30.00	22.81	32.75
Merlot	20.00	26.67	25.00	30.15	25.48
Sauvignon	36.36	17.78	33.33	15.57	25.76
Sémillon	33.33	29.63	77.78	16.96	39.43
Mean PrC per collection	41.27	18.07	53.22	20.96	

G1, first generation; G2, second generation; Pr, parasitism rate; PrC, relative contribution of the most abundant larval parasitoid.




Figure 2: Effect of different grape cultivar consumed by *Lobesia botrana* as a larva on the parasitism rate of its eggs by *Trichogramma evanescens* (N° of parasitized eggs per female of *Trichogramma evanescens*, see Moreau *et al.*, 2009 for experimental details).

These first studies strongly suggest that grape cultivar may also interfere with the host quality and thus their parasitism rate. However in the study done by Xuéreb & Thiéry (2006), the five cultivars tested did not significantly influence the larval parasitism by *C. capitator*, except Merlot and possibly Sauvignon on which *L. botrana* larvae were less parasitized by *C. capitator* in first spring generation (see Table 3).

Conclusion

A quick and simple answer to the question asked in the title should be yes. However successful use of natural enemies as a pest control techniques requires important efforts concerning the basic knowledge of the parasitoid or predator species and their relation with their targeting hosts. Also one key point in biological control is how to favour survival and high fecundity of the beneficial organisms. Research has thus a challenge to improve its knowledge of tritrophic interactions in vineyards which are to date not enough investigated. Viticulture has also to evolve towards practices that are more environmentfriendly, thus favouring the populations of natural enemies by offering them fairly favourable environmental conditions. Coupling releases of natural enemies naturally occurring in vineyards and vineyard/landscape management techniques in order to favour the installation and reproductive success of the beneficial organisms is an interesting way for future efficient biological control in vineyards.

References

- Audouin, V. 1842: Histoire des insectes nuisibles de la vigne et particulièrement de la pyrale. Fortin Masson et Cie, Paris.
- Barnay, O., Pizzol, J., Gertz, C., Kienlen, J. C., Hommay, G., & Lapchin, L. 1999: Host density-dependance of discovery and exploitation rates of egg patches of *Lobesia botrana* (Lepidoptera: Tortricidae) and *Ephestia kunhiella* (Lepidoptera: Pyralidae) by the parasitoid *Trichogramma cacoeciae* (Hymenoptera: Trichogrammatidae). J. Econ. Entomol. 92: 1311-1320.
- Barnay, O., Hommay, G., Gertz, C., Kienlen, J. C., Schubert, G., Marro, J. P., Pizzol, J., & Chavigny, P. 2001: Survey of natural populations of *Trichogramma* (Hym., Trichogrammatidae) in the vineyards of Alsace (France). J. appl. Entomol. 125: 469-477.
- Castaneda-Samoya, O. R., Holst, H. & Ohnesorge, B. 1993: Evaluation of some *Trichogramma* species with respect to biological control of *Eupoecillia ambiguella* and *Lobesia botrana* Schiff (Lepidoptera: Tortricidae). Z. Pflanz. und Pflanz. 100: 599-610.
- Chuche, J., Xuéreb, A. & Thiéry, D. 2006: Attraction of *Dibrachys cavus* (Hymenoptera, Pteromalidae) to its host frass volatiles. J. Chem. Ecol. 32: 2721- 2731.
- Collombera, S., Alma, A. & Arzone, A. 2001: Comparison between the parasitoids of *Lobesia botrana* and *Eupoecillia ambiguella* in conventional and integrated vineyards. IOBC/WPRS Bull. 24(7): 91-96.
- Coscolla, R. 1980: Aproximacion al studio del parasitismo natural sobre *Lobesia botrana* Den. Y Schiff. en las comarcas viticolas. Boll. Serv. Plaga 6: 5-15.
- Genini, M. 2000: Antagonistes de la cicadelle verte et des vers de la grappe dans le vignoble valaisan et les milieux environnants. Rev. Suisse Viti. Arbo. Hort. 32: 153-160.
- Hommay, G., Gertz, C., Kienlen, J. C., Pizzol, J., & Chavigny, P. 2002: Comparison between the control efficacy of *Trichogramma evanescens* Westwood (Hymenoptera: Trichogrammatidae) and two *Trichogramma cacoeciae* Marchal strains against grapevine moth (*Lobesia botrana* Den. & Schiff.), depending on their release density. Biocontrol Sci. Techno. 12: 569-581.

- Hommay, G., Kienlen, J. C., Gertz, C. Bihri, C. & Pizzol, J. 2011: Factors affecting the post release dispersal of *Trichogramma cacoeciae* Marchal in the vineyards. IOBC/WPRS Bull. 67: 207-214.
- Jolicoeur, H. 1894: Les ravageurs de la vigne. F. Michaud, Reims, France.
- Lane, S. D., Mills, N. J. & Getz, W. M. 1999: The effect of parasitoid fecundity and host taxons on the biological control of insect pests: the relationship between theory and data. Ecol. Entomol. 24: 181-190.
- Le Ralec, A., & Wajnberg, E. 1990: Sensory receptors of the ovipositor of *Trichogramma maidis* (Hym. Trichogrammatidae). Entomophaga 35: 293-299.
- Marchesini, E. & Della Monta, L. D. 1994: Observations on natural enemies of *Lobesia botrana* (Den. et Schiff.) (Lepidoptera, Tortricidae) in Venetian vineyards. Boll. Zool. Agr. Bachic. 26: 201-230.
- Moreau, J., Richard, A., Benrey, B., & Thiéry, D. 2009: The influence of plant cultivar of the grapevine moth *Lobesia botrana* on the life history traits of an egg parasitoid. Biol. Control 50: 117-122.
- Moreau, J., Villemant, C., Benrey, B., & Thiéry, D. 2010: Species diversity and composition of larval parasitoids of the European grapevine moth (*Lobesia botrana*): the influence of region and cultivar. Biol. control 54: 300-306.
- Maher, N., & Thiéry, D. 2006: *Daphne gnidium*, the possible native host plant of the European grapevine moth *Lobesia botrana*, stimulates its oviposition. Is a host shift relevant? Chemoecology 16: 135-144.
- Reda Abd el Monsef, A. I. 2004: Biological control of grape berry moths *Eupoecilia ambiguella* Hb. and *Lobesia botrana* Den. et Schiff. (Lepidoptera Tortricidae) by using egg parasitoids of the genus *Trichogramma*. Thesis Universität Giessen, 103 pp.
- Sentenac, G., & Thiéry, D. 2008: Les méthodes de lutte biologiques ou biotechniques contre les insectes et acariens nuisibles à la vigne. In: Conf. Int. Modiaviti, Bordeaux: 29-41.
- Stengel, M., Voegelé, J., & Lewis, J. W. 1977: *Trichogramma*. V.b. Winter survival strain and the discovery of *T. cacoeciae* Mar. on egg masses of *Ostrinia nubilalis* Hubn. in the agroclimatic conditions of Alsace. Ann. Zool. Ecol. Anim. 9: 313-317.
- Thiéry, D. 2008: Les tordeuses nuisibles à la vigne. In: Ravageurs de la vigne, Féret, Bordeaux.
- Thiéry, D. & Xuéreb, A. 2003: Relative abundance of several larval parasitoids of *Lobesia botrana* on different varieties of grapes. IOBC/WPRS Bull. 26(8): 135-139.
- Thiéry, D. & Xuéreb, A. 2004: Vers une lutte biologique contre Eudémis (*Lobesia botrana*). In: Conférence Internationale Mondiaiviti Bordeaux: 47-52.
- Thiéry, D., & Moreau, J. 2005: Relative performance of European grapevine moths (*Lobesia botrana*) on grapes and other hosts. Oecologia 143: 548-557.
- Thiéry, D., Xuéreb, A., Villemant, C., Sentenac, G., Delbac, L., & Kuntzman, P. 2001: Larval parasites of vineyards tortricids: a brief overview from 3 french vine growing areas. IOBC/WPRS Bull. 24(7): 135-142.
- Thiéry, D., Yoshida, T., & Guisset, M. 2006: *Phytomyptera nigrina* (Meigen) (Diptera, Tachinidae) parasite of the first generation of the European grapevine moth larvae in several vineyards of the Roussillon area. The Tachinid times 19: 1-4.
- Xuéreb, A., & Thiéry, D. 2006: Does natural larval parasitism of *Lobesia botrana* vary between years, generation, density of the host and vine cultivar? Bull. Entomol. Res. 96: 105-110.