



Using crop diversity to lower pesticide use: Socio-ecological approaches

Eva Thomine^{a,b,*}, John Mumford^c, Adrien Rusch^d, Nicolas Desneux^{a,**}

^a Université Côte d'Azur, INRAE, CNRS, UMR ISA, 06000 Nice, France

^b L@bisen, ISEN Yncréa Ouest, 29200 Brest, France

^c Imperial College London, Silwood Park Campus, Ascot SL5 7PY, United Kingdom


^d INRAE, ISVV, Univ. Bordeaux, Bordeaux Sciences Agro, UMR SAVE, F-33883 Villenave d'Ornon, France



HIGHLIGHTS

- Agriculture needs to be economically, socially and environmentally sustainable.
- Increasing crop diversity at the landscape scale could improve biocontrol.
- Complementary skills need to be gathered for optimal decision making.
- The landscape biocontrol management has to be established in a win-win process.
- More technical studies are needed before broadening this promising practice.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 29 April 2021

Received in revised form 31 August 2021

Accepted 1 September 2021

Available online 6 September 2021

Editor: Jay Gan

Keywords:

Landscape crop diversity

Natural enemies

Economic feasibility

Farmer acceptance

Farmer training

Farming practices

Government subsidies

Agricultural market opportunities

ABSTRACT

The farming practices adopted since the end of the Second World War, based on large areas of monocultures and chemical use, have adversely affected the health of farmers and consumers and dramatically reduced farmland biodiversity. As a consequence, many studies over more than twenty years have stated that agriculture is facing three main challenges: (1) feeding the growing world population (2) with more environmentally friendly products (3) at a reasonable return for the producer. Increasing the efficacy of biocontrol could be one lever for agriculture to meet these expectations. In this study we propose implementation of a relatively under-researched system based on the management of landscape level crop diversity that would reduce demand for pesticide use and increase conservation biocontrol. The principle of manipulating crop diversity over space and time at a landscape scale is to optimize resource continuity, such as food and shelter for natural enemies to increase biocontrol services, reduce pest outbreaks and crop losses. The feasibility of such management options is discussed in relation to environmental, social and economic aspects. The operational and institutional inputs and conditions needed to make the system work are explored, as well as the potential added values of such a system for different stakeholders.

© 2021 Elsevier B.V. All rights reserved.

* Correspondence to: E. Thomine, Université Côte d'Azur, INRAE, CNRS, UMR ISA, 06000 Nice, France.

** Corresponding author.

E-mail addresses: eva.thomine@gmail.com (E. Thomine), nicolas.desneux@inrae.fr (N. Desneux).

Contents

1.	Introduction	2
2.	Exploiting crop diversity to design pest suppressive landscapes.	3
2.1.	Potential for reducing pest pressure through landscape crop diversification	3
2.2.	Levers to promote biocontrol	4
2.3.	Choice of the scale to diversify crops	5
3.	Meeting stakeholder needs and sticking to market realism to apply crop diversification	5
3.1.	Socio-economic aspects – how to get stakeholder acceptance for such systems?	5
3.1.1.	Questions about farmers technical capacity, added value of the system and the right business model to adopt	6
3.1.2.	Implementing decisions at a large scale	7
3.1.3.	The importance of establishing the system in a win-win process.	8
3.1.4.	Potential difficulties that could be encountered for building-up a landscape system based on crop diversity and conservation biological control.	8
3.2.	Actions to be taken by the stakeholders to reach the next step.	9
	Declaration of competing interest.	10
	Acknowledgement	10
	References	10

1. Introduction

Intensive agriculture has negative effects on the environment and on human health (Tschardt et al., 2005; Jokanovic, 2018; Forkuoh et al., 2018; Wahlang, 2018). These negative effects are caused by habitat destruction, low crop diversity, intensive soil tillage and intensive use of agrochemicals, including pesticides and fertilizers (FAO, 2019). Pesticides have been used in agriculture for over a century to increase food production and they have proven their efficiency in increasing food accessibility worldwide (Pingali, 2012; Nelson and Burchfield, 2021). Now agriculture is facing the negative consequences of the intensive use of pesticides, notably through the increase of health issues affecting farmers (Jokanovic, 2018) and consumers (Forkuoh et al., 2018; Wahlang, 2018), and also through the destruction of biodiversity in fields and surrounding land (Liu et al., 2018), on soil biota (Lew et al., 2009; Velki et al., 2019), water (Leach and Mumford, 2008; Ibrahim et al., 2019), and on arthropod biodiversity including natural enemies (Desneux et al., 2007; Van der Valk et al., 2011; Lundgren et al., 2013; Woodcock et al., 2017; Wagner, 2020). These natural enemies can be microscopic (fungi, bacteria, virus and nematodes) (Lacey et al., 2001) and macroscopic (predators and parasitoids) (Stiling and Cornelissen, 2005). In addition to negative effects due to pesticides, the use of large scale monocultures makes it difficult for natural enemies to find food and shelter after the crop is harvested, resulting in the loss of their populations and a reduction in biological control impact (Schellhorn et al., 2014). Biological control consists of “the use of living organisms - i.e. natural enemies - to suppress the population density or impact of a specific pest organism, making it less abundant or less damaging than it would otherwise be” (Eilenberg et al., 2001). Changing pest management practices by increasing the biological control potential, including the reduction of pesticide use, is an objective for the future as agriculture faces three challenges: to sustain healthy food production for the growing world population, to reduce the negative impacts of agrochemicals on the environment and on human health, and ensure reasonable profit or return for the producer.

Natural enemies can be used in agriculture in several ways (inundative, classical and conservation biological control) (Bale et al., 2008). Conservation biological control using macroscopic natural enemies, on which we will focus in this study, works by managing the environment to promote naturally occurring natural enemies (Eilenberg et al., 2001). Its main principle is to “enhance the activity of existing natural enemies to provide pest suppression” (Haan et al., 2021) notably by increasing plant diversity (Andow, 1991; Altieri, 1999) providing continuous access to diversified food sources (pollen, nectar, alternative hosts and prey) and shelter, despite harvest, crop senescence, or even

pesticide use in some fields (Jonsson et al., 2008; Gurr et al., 2017), as well as overwintering sites (Gurr et al., 2017; Haan et al., 2021) between seasons.

Increasing plant, i.e. crop or non-crop, species diversity (referred to as inter-specific diversity later in the paper) (Andow, 1991; Letourneau et al., 2011; Wratten et al., 2012; Nicholls and Altieri, 2013), plant genotypic and phenotypic diversity (referred to as intra-specific diversity later in the paper) (Koricheva and Hayes, 2018) and plant functional diversity (Hatt et al., 2020) across spatial and temporal scales has been found to benefit biological pest control services and limit yield losses by increasing the presence and the activity of natural enemies (top-down effects) and by reducing pest pressure (bottom-up effects). Plant diversification includes both crop and non-crop habitats and can be managed at the field (e.g. flower and grass strips, intercropping), farm (e.g. crop rotations) and landscape scales (e.g. hedgerows, forest) (Letourneau et al., 2011; Lin, 2011; Jeanneret et al., 2012). During the last thirty years of research a large number of studies focused on inter-specific diversity at the field scale or on non-crop habitat density at the farm and landscape scale and demonstrated beneficial effects of these diversification schemes on biological pest control services (Bianchi et al., 2006; Rusch et al., 2016; Karp et al., 2018). However, few studies investigated how crop diversification (within and between crop species) at the landscape scale could be a major management option to enhance biological pest control services in agricultural landscapes. Diversifying crop species in space and time can not only be positive for ecosystem services but might also be positive from economic and social aspects (Craheix et al., 2016).

In this paper, we envision that cropping patterns with intra-specific and inter-specific diversity at the landscape scale might be a key management options to promote biological pest control services and to ensure greater and more stable incomes for farmers while limiting negative externalities related to farming activities (Nicholson and Williams, 2021). We first address how crop diversification at the landscape-scale could be a major management option to limit pest pressure and define the ecological requirements to optimize biological control of pests in agricultural landscapes. Then, we propose ways to meet socio-economic needs of stakeholders, conditions for acceptance of such innovations and technical opportunities to overcome difficulties in applying such management options in real-life landscapes.

In the rest of the paper we will refer to an agricultural system, which is a system where crops are diversified, intra or inter-specifically, at a landscape scale. The landscapes are defined as areas shared between humans, flora and fauna, considered at different radial dimensions depending on the species observed – a 1.5 km radial unit is usually




Fig. 1. Schematic representation of two contrasted sites with low crop richness (A) and high crop richness (B). Water is represented by blue, natural habitats by dark green and human-mediated uses by dark grey. All the other land uses are different types of crops.

appropriate to achieve an effect from landscapes on arthropods (Gardiner and Neal, 2009) – and often contains different types of land uses, such as urban areas, natural areas and cropped areas (see Fig. 1 for more details on the landscape considered). The socio-economic network studied in this paper is composed of farmers, environmentalists, retailers, consumers and policy makers. These actors are all linked by diversified relationships and are connected to environmental, social and economic pillars through a range of individual purposes, objectives and hopes (see Fig. 2). For example, policy makers hope to sustain the State economy by creating new projects and at the same time, aim at sustaining social stability with diverse financial supports. Policy makers also have the global objective to protect the environment by creating laws forbidding or, conversely, encouraging some farming practices - for example, greening of the Common Agricultural Policy (CAP)

proposed by the EU Member States (Matthews, 2013). Farmers, on the other hand, aim to maintain sustainable markets in order to preserve the economy of their enterprises and at the same time aim to work with environmentally friendly techniques in order to preserve their health and the health of consumers. Finally, an indirect farming objective, often not explicitly claimed, is to preserve the environment as the farming industry is in permanent interaction with it.

2. Exploiting crop diversity to design pest suppressive landscapes

2.1. Potential for reducing pest pressure through landscape crop diversification

Crop diversification across spatial and temporal scales can affect pest populations dynamics through two non-exclusive mechanisms:




Fig. 2. Schematic representation of the agricultural network studied in relation to the economic, social and environmental pillars. In grey are represented the stakeholders targeted, in capital letters and bold arrows are the different relations between stakeholders. The thin arrows represent purposes (normal arrows), objectives (dotted arrow) and hopes (dashed arrow) of each stakeholder toward each pillar.

bottom-up (the resource concentration hypothesis) and top-down (the natural enemy hypothesis) effects. On one hand, bottom-up effects can be activated by diluting the plant resources used by pests. Diversifying plant species and/or genotypes has been demonstrated to be efficient in reducing pest pressure as individuals are less able to find their food sources across the agricultural fields (Letourneau et al., 2011; Koricheva and Hayes, 2018; Snyder et al., 2021; Wan et al., 2020; Li et al., 2020). In a recent review Koricheva and Hayes (2018) have highlighted that crop genotype diversity seems to have a stronger effect in reducing pest pressure than does wild plant genotype diversity. The authors explain this difference as an effect of associational resistance to pests being stronger (Root, 1973) against a specifically targeted pest in crop experiments than in wild plant experiments (Koricheva and Hayes, 2018). On the other hand, top-down effects can be activated by increasing the accessibility of diversified food resources and shelter to natural enemies in order to enhance their abundance and performances (Letourneau et al., 2011; He et al., 2019). To date, both inter-specific and intra-specific diversity has demonstrated positive effects on natural enemies and pest reduction but only crop inter-specific diversity effects on arthropods have been studied at the landscape scale, studies on the effect of crop intra-specific diversity being limited to single field studies (Koricheva and Hayes, 2018; Snyder et al., 2021). We therefore detail in the following section only the effects of crop inter-specific diversity at a landscape scale on natural enemies, pests and biocontrol (i.e. predation and parasitism).

A growing number of studies have addressed the question of landscape inter-specific crop diversity impact on natural enemies, pests and biocontrol (Bosem et al., 2017; Liu et al., 2016, 2018; Redlich et al., 2018; Sirami et al., 2019; Aguilera et al., 2020; Kheirodin et al., 2020; Zhao et al., 2021). These studies related various effects of agricultural landscape crop diversity: (1) positive effects on parasitism (Liu et al., 2016), predation (Redlich et al., 2018), on the abundance of natural enemies (Liu et al., 2018; Zhao et al., 2021) and their diversity (Aguilera et al., 2020; Sirami et al., 2019; Zhao et al., 2021), (2) negative effects on the abundance of pests when crops were not host crops (Bosem et al., 2017; Kheirodin et al., 2020) (support of the resource concentration hypothesis) and (3) a higher ratio of natural enemies to pests (Zhao et al., 2021). Diversifying agricultural landscapes through crop manipulation can therefore have a positive impact on biodiversity of natural enemies, even higher than from semi-natural habitats (Sirami et al., 2019), and on biocontrol, when many different crop types are grown or when crop hosts are not usually cultivated in the same landscape unit of potential interaction. Even though the number of papers reporting the effect of landscape crop inter-specific diversification is modest, we can presume that using crop inter-specific diversity at a large scale might be positive for natural enemies and/or on biocontrol as suggested by Nicholson and Williams (2021) or Larsen and Noack (2021).

The positive effects of crop inter-specific diversity at a landscape scale can be the consequence of two main mechanisms: 1) complementary landscape hypothesis for natural enemies enhancement (top down control of pests) and 2) resource dilution hypothesis for pest reduction (bottom-up control of pests). Concerning the complementary landscape hypothesis, alternative crops can act as reservoirs for natural enemies that can spill over from one crop type to another when resources are increasing, notably pest resources (Liu et al., 2018). They can also act as shelter and food resources when the main crops are harvested or treated with pesticides (Liu et al., 2018; Aguilera et al., 2020). Parasitism rates can be increased through direct bottom-up forces: the increase of crop species can increase the presence of generalist pests and in consequence can increase the parasitism rates on this pest by parasitoids that can find their main host in multiple crops (Liu et al., 2016). Some authors underline the importance of combining both crop and semi-natural habitat diversity at a landscape scale for more biocontrol efficacy, defined as potential to reduce pest abundance (the complementary landscape hypothesis

suggests a strong complementarity of semi-natural habitats and crops for resources and shelter) (Sirami et al., 2019; Aguilera et al., 2020). Sirami et al. (2019) found that the proportion of semi-natural area cover in a landscape had a positive effect on the level of increase of multi-trophic arthropod diversity as crop species diversity was increased in the landscape. They show that crop diversity is particularly important in maintaining arthropod diversity when the proportion of semi-natural cover in the landscape is very low. The authors suggest that complementarity of both crop and semi-natural areas comes from spatial and temporal resource continuity given by crop inter-specific diversity and shelter given by semi-natural areas (Sirami et al., 2019). More than crop diversity, choice of crop types to include can highly influence the diversity and abundance of natural enemies, as has been shown in oilseed rape for spiders and carabids (Aguilera et al., 2020) suggesting that only increasing inter-specific diversity might not be sufficient, but paying attention to the function of the specific crop types involved should also be part of the decision. Finally, diversifying crop types at a landscape scale increases natural enemy community diversity, which implies higher potential for pest control (Zhao et al., 2021). Concerning the resource dilution hypothesis, increasing crop inter-specific diversity has been demonstrated to reduce pest abundance, more specifically specialist pest abundance, reducing their capacity to find their principal host plant (Bosem et al., 2017; Kheirodin et al., 2020). It is important to mention that even though proofs of the potential high effect of crop intra-specific diversification, through genetic diversification of crops, at a landscape scale are absent from the literature. This lack of studies on the intra-specific diversity effects at the landscape scale seems to come from the complexity of the potential interactions and the crop species quality issue. There are a lot of potential types of diversity, but some have more diverse quality than others and so the effects are very variable. This species quality issue needs to be studied further as it shows substantial potential in pest reduction and natural enemies increase.

2.2. Levers to promote biocontrol

Diversifying crops at a landscape scale appears to have promising potential to reduce pest pressure by increasing natural enemies presence and diversity in arable fields. In order to maintain pest populations at a low level, it is important to maintain diversified guilds of natural enemies with complementary resources needs, i.e. different guilds of pests, in fields (Symondson et al., 2002). Natural enemies have different needs in terms of resources (i.e. pollen, nectar, or alternative preys and hosts) and habitats to realize their life-cycle (e.g., overwintering sites, shelter during the summer) (Gurr et al., 2017; Gardarin et al., 2018; Symondson et al., 2002). Natural enemies can be specialists (e.g. most parasitoids that specifically attack specific species of aphids) (Fischbein et al., 2016; Monticelli et al., 2019) or generalists (e.g. spiders, ladybirds or carabids that can feed on different types of pests) (Rand and Tschardtke, 2007) and can have specific needs at different stages during their development (e.g. only proteins) or various needs (e.g. pollen and nectar during the adult stage and proteins during the larval stage, as is the case of most parasitoids and hoverflies) (Van Rijn et al., 2013; Fischbein et al., 2016). Positive relationships between species richness of natural enemies and pest suppression have been reported and emerge due to niche partitioning or sampling effects (Letourneau et al., 2009). Maintaining diverse communities of natural enemies is therefore an important lever to efficiently control different pest population types in space and time (Dainese et al., 2017).

Understanding the movement of natural enemies is another key to design pest-suppressive landscapes. The movement of natural enemies can be driven by multiple factors: biotic factors – such as the presence of conspecifics (Tuda and Shima, 2002), herbivore-induced plant volatiles (Gillespie et al., 2016), plant phenological stages (Schellhorn et al., 2014), movement capacity of the guild (Osawa, 2000; Chapman et al., 2006; Jauker et al., 2009; Wang and Keller, 2003) – and abiotic factors – such as the climate (Schellhorn et al., 2014). However, movements

of arthropods in a landscape are also conditioned by landscape structure, both in terms of composition and configuration (Karp et al., 2018; Martin et al., 2019; Haan et al., 2020). The amount of habitat sources for natural enemies, such as semi-natural habitats, is the main determinant of natural enemy presence and abundance (Sirami et al., 2019). It has been recently suggested that a minimum of 20% of semi-natural habitats within a few kilometers, such as forests or natural grasslands, is needed to maintain a significant pool of natural enemy species (Tscharnkte et al., 2007; Garibaldi et al., 2021). In addition to the amount of habitats, the spatial configuration of habitats in the landscape affects movements of natural enemies. Natural enemies are usually more abundant in fine-grained agricultural landscapes, i.e. landscape with smaller patches of habitats, that enhance spillover of natural enemies and connectivity (Bailey et al., 2010; Martin et al., 2016; Haan et al., 2020). Reducing the amount of roads, hedgerows and tree lines that can act as shelter and increasing edges between crops and corridors are criteria to take into account to optimize biocontrol in agricultural landscape (Schellhorn et al., 2014). However, it should also be noted that for some natural enemies these landscape features may act as barriers to movement, rather than enhancing access. Evidently the relative importance of landscape composition and configuration for natural enemies depends on specific natural enemy traits, such as dispersal ability (Martin et al., 2019; Haan et al., 2020).

Landscape compositional and configurational traits, as well as arthropod needs and capacity to disperse, are important factors to consider in order to maximize our chances to reach a landscape rich in biocontrol potential (Haan et al., 2021). Spatial and temporal resource continuity is a key to increase natural enemy spillover from one pest host resource to another with no decrease in their population abundance (Vasseur et al., 2013; Schellhorn et al., 2015; Iuliano and Gratton, 2020). Selecting crops to reach a successful diversification scheme should therefore consider: 1) the landscape crop composition, such as the complementarity of resources given by the crops available for a high number of natural enemies (pollen and nectar provision but also alternative preys and potential shelter) (Gardarin et al., 2018); 2) the landscape crop configuration (Haan et al., 2020), such as fields size, field shape, field connectivity making the resources continuous in time and space by selecting smaller crop areas that follow and overlap each other in time to maintain a continuous food source and shelter for natural enemies (Vasseur et al., 2013; Schellhorn et al., 2015; Sirami et al., 2019; Nicholson and Williams, 2021). We do not refer to any specific species associations, as conditions in each location can change regarding climate, soil, landscape composition, etc.

2.3. Choice of the scale to diversify crops

Diversifying crops in order to suppress pests and/or increase natural enemy efficacy can be done at three different scales: (1) at the field scale – generally through polyculture schemes (companion cropping, push-pull, intercropping, trap-crops, etc.) (Letourneau et al., 2011; Beillouin et al., 2019) or through rotation schemes (Rusch et al., 2013; Barzman et al., 2015; Beillouin et al., 2019) activating bottom-up forces through the resource dilution hypothesis, (2) at the farm scale – diversifying crops in multiple fields but in one farm only (Jeanneret et al., 2012), and (3) at the landscape scale – a scale that has been only recently studied as shown in Section 2.1 (see Fig. 3 for more details on the different scales described). Manipulating plant diversity at the field scale seems to be most efficient, but for issues of technical feasibility these systems are under-used in modern agriculture (Schaller, 2012; Meynard et al., 2013; Meynard et al., 2018; Morel et al., 2020). In order to avoid any conflicting effect of practices on the efficacy of natural enemies (Brittain et al., 2010) applying the proposed agricultural system based on crop diversity and reduced pesticide use at a large scale would be the most efficient scale (Landis, 2017; Brewer and Goodell, 2012; Goldman et al., 2007; Haan et al., 2021). Additionally, as argued by Landis (2017), even though a particular farm is efficient, in terms of

biocontrol increase through crop diversification, other less efficient farms interspersed with the efficient farm might reduce the overall efficacy of the method used by that efficient farmer (Landis, 2017), for example in the case of pesticide drift. Taking decisions at larger scales than individual farms is therefore a key to success but will require efficient planning and coordination among different farms (Landis, 2017; Haan et al., 2021).

Arthropods, especially large ones (e.g. ladybirds, lacewings, hoverflies, but also spiders), can move over long distances (ranging up to several kilometers) in order to find their food and mate (Roh, 2013; Evans, 2003; Chifflet et al., 2011; Villenave-Chasset, 2006). Studying the effect of a cropping system on a small or partial landscape, such as a field or farm, would therefore omit a large part of the landscape covered by the natural enemies, and consequently the impact of the rest of the landscape on these animals. In landscape ecology, the landscape scale to study a broad spectrum of natural enemies is usually between 1 km (e.g. Rusch et al., 2016) and 2 km radius (e.g. Karp et al., 2018). The study of Gardiner and Neal (2009) has shown that the 1.5 km scale is best to explain the variation in biocontrol and abundance of ladybirds, a large long distance flying predator. Many levers of landscape manipulation for natural enemy preservation have been shown to be effective (Landis, 2017): 1) landscape heterogeneity needs to be preserved and both composition and configuration of the landscape, not only composition as developed in the previous section, need to be considered when managing a landscape (Holzschuh et al., 2010; Fahrig et al., 2011; Perovic et al., 2015), 2) landscapes need to be connected and field sizes reduced in order to allow spillover between fields and entire exploitation of resources in the fields (Fahrig et al., 2015; Fischer et al., 2006; Haan et al., 2020), 3) food provision or natural enemies needs to be continuous in time and space (Schellhorn et al., 2015) and 4) disturbing events, such as ploughing, harvesting, vegetation clearance, cutting and pesticide treatments, need to be adapted to arthropod life cycles (Fischer et al., 2013).

If biocontrol is to be increased through crop diversification, large monoculture fields might need to be divided into multiple small fields of different crop types. Dividing large fields into long narrow fields of effective polyculture might facilitate natural enemy circulation between fields of different crops and at the same time decrease the possibility for pests to find their host plants (resource dilution hypothesis) and facilitate crop management by farmers. Indeed, this way of arranging fields has been applied for a long time in China and has been proved, when applied at a landscape scale, to be efficient in increasing the abundance and species richness of natural enemies in the cultivated fields (Zhao et al., 2021). These technical decisions on field shape will therefore need to match 1) ecological needs of natural enemies in terms of movement, wind direction, etc., in order to optimize biocontrol and 2) farm management needs for a simple system to manage.

3. Meeting stakeholder needs and sticking to market realism to apply crop diversification

3.1. Socio-economic aspects – how to get stakeholder acceptance for such systems?

Even though the willingness of farmers to switch from intensive agriculture to more environmentally friendly techniques is increasing, especially with government agri-environmental schemes offering incentives (Bernués et al., 2016; Wezel et al., 2018), the long-term application of pesticides has locked farmers into a situation where it is economically risky to switch to a zero-pesticide system; this situation is called the socio-technical lock-in (Wilson and Tisdell, 2001; Guichard et al., 2017; Meynard et al., 2018), where fear of losing economic gains and uncertainty of new techniques combine to dissuade farmers from switching to another system (Guichard et al., 2017). We propose here to meet the three main objectives needed to achieve a change in agricultural practices. One objective is to address the




Fig. 3. Representation of the different scales where crop diversity and biocontrol can be managed. The regional scale is a scale defined politically where decisions can be taken broadly. The landscape scale is a scale defined by landscape ecologists (at least in our case study) where arthropods movements and behavior can be observed and linked to an agricultural practice. The farm scale is a scale defined economically by farmers where management decisions can easily be taken by individuals. The field scale is a scale defined economically by farmers where tests can be done without implying too much economical risks.

questions to ask in order to enter the new system. Another objective is to determine the spatial and temporal scale at which this system might be best applied. Finally, a last objective is to consider the importance of establishing this system as a win-win process to achieve rapid and sustainable adoption. We then pose three main challenges that could be encountered when trying to establish a large-scale crop diversification scheme to increase biocontrol and reduce pesticide uses.

3.1.1. Questions about farmers technical capacity, added value of the system and the right business model to adopt

After understanding clearly the problems articulated by farmers (economic, social and environmental issues), the first question to address is the way to efficiently train farmers the role of crop diversity (within their overall farming objectives) in order to boost natural enemy activity and increase economic efficiency. Modern farming is first of all an enterprise where profitability is a major objective (Bernués et al., 2016). Responding to farmer needs through training toward new environmentally friendly concepts is often associated with ideas which are not seen to have immediate direct and obvious economic return for the farmer (Kilpatrick and Rosenblatt, 1998). Above all, it is necessary that, when learning together as farmers and ecologists, farmers gain knowledge relevant to their personal situation, by demonstrating concepts associated with practical examples (Kilpatrick and Rosenblatt, 1998). Finally, training farmers must 1) overcome any lack of confidence farmers may have in trainers when training is compulsory and in the accuracy of new information given, 2) overcome

any fear of learning new knowledge or skills as it might induce a change of practices and habits, problems that have been found to be the main reasons why training fails to reach or be adopted by farmers (Kilpatrick and Rosenblatt, 1998) and 3) show that working together as farmers and ecologists to develop a training curriculum is relevant to the needs of growers and opens up new practical and effective management options for them. Implementing a new agronomic system will therefore need to be done with a clear view of the added value for the farm owner. One additional advice would be for other farmers who have already tested the system to train further farmers to ensure the efficiency of the program proposed and to increase the relevance of training (Kip-Tot et al., 2011; Bouttes et al., 2019). Local actions led by community organizations and group training could be important, to involve farmers directly in the decisions and the organization of the landscape (Stallman and James, 2015). Landscape re-organization in accordance with farmer needs (social and economic) and with the ecological intensification of agriculture will likely need some public coordination and State finance (Fahrig et al., 2011). Implementing a self-learning/self-training program in which farmers convince themselves of good practices would also be useful to encourage farmer groups to experiment at a local scale with landscape coordination and keep records of impacts on natural enemies and pest challenge (affecting pesticide use).

A second question that must be raised is how the farmer can get any value from the system. The potential reduction of pesticide use in a system where natural enemies could limit pest outbreaks will reduce

exposure of farmers to chemicals and will reduce the probability of pesticide related health problems (Jokanovic, 2018). Additionally, by diversifying crops, farmers might be less subjected to commodity price fluctuations (Olsson, 1988; Gilbert and Morgan, 2010; Haile et al., 2017). Crop price volatility can be due to different factors on which farmers have almost no control: rapid economic growth of some developing countries, decades of underinvestment in agriculture, poor harvests due to climate changes, currency depreciations, diversion of food crops into the production of biofuels and speculative influences (Gilbert and Morgan, 2010). Diversifying crops might reduce the pressure of these previously cited factors, which are principally acting on crops like rice and wheat (Gilbert and Morgan, 2010). Finally, constant exposure of crops to newly introduced pests due to climate change (Ziska et al., 2011; Barzman et al., 2015) and globalization (Hulme, 2003; Peña, 2013) makes a system based on diversified crops less vulnerable to yield losses (Lin, 2011; Degani et al., 2019). Agroecosystems with diversified traits and functions are more resilient to changing biotic and abiotic conditions (Lin, 2011) due to two main reasons: the system is more able to suppress pest outbreaks and pathogen transmission through the resource dilution hypotheses (Otway et al., 2005) and it can buffer the effect of climate variability on production (Lin, 2011).

Agricultural landscapes can be associated with specific markets. Agri-tourism can be an additional market to use by farmers in order to make a profit from eco-friendly agrosystems, in some situations: 1) the farm needs to be near a touristic area (Sharpley and Vass, 2006), 2) the project needs to be eligible for environmental public subsidies (Haan et al., 2021), 3) may require training on marketing techniques (Sharpley and Vass, 2006) and means of communication (websites, social networks, etc.). Another market could be local sale of newly introduced crops under an environmentally friendly label at higher prices since consumers are willing to pay higher prices for such products (Elkington, 1994; Cranfield and Magnusson, 2003). Labelling can be an option to help consumers learn about the effort made by farmers, and would allow farmers to sell products at higher prices if consumers are willing to pay more for healthier products, as shown by a survey led in the framework of the EUCLID H2020 program asking if consumers were willing to pay more for environmentally friendly – not organic – products (up to 20% more), especially for fruits and vegetables (Fornetti, 2019). However, a communication effort about these new brands will need to be done by retailers as consumers are more confident about well-established organic and fair trade labels rather than new labels (Sirieix et al., 2013). The newly introduced crops might also be sold in local markets at higher prices if the farm is located in a peri-urban place, as many consumers today prefer to choose local IPM over non-local organic (Adams and Salois, 2010; Fornetti, 2019). In case none of the markets proposed above are applicable, public subsidies might help farmers to apply environmentally friendly techniques, as for example the current CAP greening, or the Whole Farm Revenue Protection introduced in 2014 in the US allowing farmers to diversify their production in order for them to increase their resilience (Haan et al., 2021). Several systems of payment exist for subsidies based on

different units: the simple input area per hectare, the output volume per ton, the output value per currency unit, the action and action avoided payment per unit of approved or proscribed input respectively, and the outcome payment (Table 1). Regarding the different systems of payments, the action payment system would be best adapted in the agro-ecosystem considered in this study. Contrary to output payments, the objective of the system based on multiple crop farming is not to produce more but to produce better. Finally, action payments, as opposed to action avoidance payments, are morally more rewarding.

The last question to ask is how to build a business model for entry and maintenance of the system in a specific chosen market. The products delivered in a farming system where crops are diversified and produced under reduced use of pesticides are healthier and better quality, and can be sold as such. Before introducing new crops, farmers need to assess potential markets for the newly introduced crops. Retailers promoting environmentally friendly products might be the best stakeholders to target for product sales. The “Zero pesticides” tomato from the French Saveol enterprise is a good example showing that large retailers, because of an increasing demand from consumers and thanks to an adapted branding from the firms, are buying and selling more and more environmentally friendly products and therefore are good target markets (Raynaud et al., 2009).

3.1.2. Implementing decisions at a large scale

As mentioned in Section 2.3, in order to avoid any conflicting effect of practices on the efficacy of natural enemies (Brittain et al., 2010) applying an agro-ecosystem based on crop diversity and reduced pesticide use at a large scale would be most efficient (Landis, 2017; Brewer and Goodell, 2012; Goldman et al., 2007) from an ecological point of view. If we focus now on the political/social point of view, such global decisions could be difficult to organize and might take a long time before being efficiently applied at a national scale. Diversifying crops could be done first at a regional or local scale (Cumming and Spiesman, 2006; Valbuena et al., 2010). One example of agri-environmental legislation that has been implemented at a regional scale is the French Regional Action Program (RAP) initiated in Haut-de-France in 2018 for protection of water bodies. RAP is experimenting with innovative farm techniques involving nitrogen fertilizer management in order to reduce pollution by nitrates of agricultural origin in the region. Targeting smaller levels of action, such as the farm level, is also possible but might be less efficient in case farm parcels are highly interspersed with other parcels and the farming practices of other farmers are deleterious to natural enemies (Landis, 2017; Slotterback et al., 2016). At a higher level, it is possible that a group of farmers or farm unions could take the decision to increase their crop diversity and decrease pesticide use to promote natural enemies, with possible optimal biocontrol if parcels are adjacent. The design of new agricultural landscapes needs to be done through collaborative networks of different specialists (Landis, 2017; Haan et al., 2021). As stated by Landis (2017) and Haan et al. (2021), the use of different knowledge in order to answer farmer needs could be done through the mobilization of environmentalists (to understand the

Table 1

Table relating the different systems of payments proposed by the European states for agricultural subsidies. The payment name, the associated units and definition and the references are reported in the table.

Payment name	Unit	Principle	Reference
Simple input area	per hectare	farmers are offered an amount of money per hectare	Baylis et al., 2008
Output volume	per ton	subsidies support production by giving money per quantity of agricultural commodity produced	Van Zanten et al., 2014
Output value	per euro	subsidies support production by giving money per quantity of benefit produced	
Action payment	per unit of approved input	farmers can be paid to stay on their farm in order to preserve the farmland or can receive a compensation for not attaining quotas	Baylis et al., 2008
Action avoided payment	per unit of proscribed input stopped	for example when a farmer is paid for reducing chemical input or animal units per land area	Baylis et al., 2008
Outcome payments	related to water quality, residue limits, landscape assessment, etc.	payments can be received when farmers measures are taken by the farmer to protect the environment	Baylis et al., 2008; Van Zanten et al., 2014

species needs and biodiversity conservation techniques), geographers (organization of the landscape), economists (establishment of a working business plan for farmers), sociologists (understanding the social objectives and opportunities), agronomists and farmers themselves (technical input, establishment of a working technical program in the region, transfer of the techniques to other farmers). Advice during cropping periods about timing of pesticide applications and natural enemy dynamics will also be needed for maximum efficiency, such as maintenance of pest pressure under the economic threshold, and preservation of natural enemies. Indeed, a clear understanding of natural enemy dynamics and pest outbreaks will be needed to apply pesticides only when natural enemies are in low numbers, and applications may be limited to the center of fields where the natural enemy density might be lowest (Bortolotto et al., 2016). Creation of decision tools adapted to landscape scale management could be used in order to coordinate the choice of crops by different farmers regarding the population dynamics of the different insects. Such a tool has already been developed by Slotterback et al. (2016) where farmers' decisions in a region were transferred into a tool called Geodesign. An iterative process helped to assess the resulting changes happening at the landscape scale and helped in the emergence of multifunctional solutions (Slotterback et al., 2016). Networks gathering different agricultural stakeholders have also been developed all across Europe in order to help design efficient agroecological farming systems (e.g. Agroecology Europe Forum which has gathered more than 300 participants, notably farmers, technicians, researchers, students, policy and decision-makers, representatives of national and European institutions, non-governmental organizations, social movements, and civil society (Wezel et al., 2018)).

3.1.3. The importance of establishing the system in a win-win process

In the context of market greening, often initiated by legislative requirements, companies have become more competitive and innovative, benefiting in a win-win process from consumer demand for greener products (Elkington, 1994; Peattie, 2001). This win-win process is frequent in organic agricultural markets, as has been shown, for example, in a Globe Newswire interview where organic farmers in the United States were benefiting from premium prices given under an organic brand trusted by consumers (Global Newswire, 2019). Another example of a win-win process, established within a crop diversification scheme in organic crop rotations, is the brand Annie's from General Mills that is buying crops newly introduced in a rotation with previously established crops and that were previously not grown (Crawford, 2019). In an agro-ecosystem based on crop diversification and low pesticide inputs, we identified five main stakeholders potentially positively impacted by the system.

1. **First of all, farmers:** Reducing pesticide use that is bad for their health (Jokanovic, 2018) would be a high benefit for them. Increasing crop diversity could also raise new markets and push agri-food companies to buy new crops at high prices, at least during the transition phase, in order to meet consumer demand for healthier food (Crawford, 2019). The satisfaction induced by adoption of environmentally friendly methods would also improve farmer well-being (Fischer, 1980) by increasing the level of working conditions (Shreck et al., 2006). Direct help from the State in order to switch to more environmentally friendly system could help to ensure stable incomes.
2. **Then, environmentalists:** Arthropod biodiversity preservation, among other animal diversity preservation like birds and mammals, as well as the moderated environmental impacts that might result from adoption of diversified crop systems and reduced pesticide use (Letourneau et al., 2011) are clear positive arguments for environmentalists.
3. **Consumers:** Demand for healthier and environmentally friendly products would also be a "winner" in the proposed system in two ways: 1) reduction of pesticides would generate better quality

products with reduced residues (Smith-Spangler et al., 2012), and 2) reduced use of pesticides would reduce potential health problems related to drift (Provost et al., 2007).

4. **Retailers:** With an increase in consumer demand for healthy products, and increasing conversion of farmers to IPM or organic farming, retailers can be included as key actors in promotion and distribution of healthy products, under marketing processes that help to increase sale prices and therefore benefits (Crawford, 2019).
5. **Policy makers:** If the increase of crop diversity at a landscape scale can help reduce the use of pesticides, as the main objective of policy makers is to maintain public health at a high level and preserve biodiversity, this solution might also be a winning solution for them.

At this stage it is important to note that reducing the dependency of farmers on pesticides might not be a winning solution for agrochemical companies (Clapp, 2021). However, today these companies are incorporating IPM concepts through the adoption of new technologies like RNA-based biocontrol products (Taning et al., 2020), precision agriculture (Birner et al., 2021) or even biocontrol (see "Biologicals by Bayer" as an example) and might, by necessity, switch completely to IPM solutions in order to satisfy consumer and policy demands (but see Deguine et al., 2021). It is important to remember that consumption is the basis of a market, and if the demand for conventional food is reducing, the companies will have to adapt to what consumers want. More details about the influence of agrochemical firms in decision making are given in the following sub-section.

3.1.4. Potential difficulties that could be encountered for building-up a landscape system based on crop diversity and conservation biological control

We mentioned in the previous section that if crops are diversified inter-specifically, then farmers will need to find new markets in order to sell their newly introduced crop species in their cropping system. Creating new markets might not always be easy if the demand is not present. However, one way to overcome that difficulty would be to diversify the cropping systems intra-specifically, by diversifying the genetics of a cultivated species. Choosing this option will avoid the difficulty of finding new markets in case there is no demand for the proposed new crops (Koricheva and Hayes, 2018).

A second possible blocking point might be the feasibility of implementing the proposed landscape design at a large scale. Communication between farmers to preserve natural enemy communities in fields is necessary as agricultural landscapes are composed of different farms (Cumming and Spiesman, 2006; Goldman et al., 2007; Stallman and James, 2015). Coordination of practices applied by different farmers of a region (Stallman and James, 2015) might help to optimize choice of crops to implement, pesticide use and resulting biocontrol services. Cooperation between farmers on practices to increase biocontrol is possible but might not always work, as farmers with more inclusion in community organizations or farmers that are concerned about pesticides in the environment seem more willing to cooperate than those who don't (Stallman and James, 2015). Local scale (neighboring farms) cooperation might also be more efficient than a larger county-wide scale (Stallman and James, 2015). All in all, cooperation between farmers, that could be enhanced by regional managers, might be possible only at small scales where farmers agree on principles of pesticide reduction and timing of applications that allow natural enemies to establish in the landscape. A last point where farmers may need to collaborate in a diversified crop landscape is on purchase of agricultural equipment. The management of different crop types requires different type of equipment: combines for cereals, mowers for fodder, harvesters and leaf strippers for beets, but also various types of seed drills at the beginning of the cropping season, as well as specific cultivation, etc. Increasing the number of crops on a farm will require more machine types and might be very expensive if farmers must purchase these machines alone. To share purchase of equipment, as is already done in

France with the CUMAs (Cooperatives for the Use of Agricultural Machineries), might be a good option.

A third blocking point might be that the proposed landscape design might not be easy to implement in all types of farms. Even though we mainly specified that landscape crop diversity needs to be managed at a landscape scale, it is possible that some farms might have more or less difficulties to implement such management practices. The size of the farm might be one excluding criteria. The size of a farm can be defined in relation to its area cultivated or its capital (standard gross margins) (Potter and Lobley, 1993; Nagayets, 2005). Based on the economic status of each type of farm, diversifying crop systems may be more easily adopted by small farms (35 ha large and less) (Burton et al., 1999; Rigby et al., 2001) at the season scale through rotations. As small scale farmers have generally little capital investment, they may have more flexibility through contracted equipment suppliers and therefore may be more able to switch easily from one crop type to another from season to season. However, implementation of diversifying crop practices and lower pesticide inputs to increase biocontrol might have more impact in areas where large scale monocultures are usually farmed. Large scale farms (above 100 ha, see Burton et al., 1999) could more easily diversify their crops in space and time as they usually have more capital (Haspel, 2014) and have more area to work on. The larger the farm, the greater the investments can be and the production cost per unit goes down (Haspel, 2014). We raise the hypothesis that risks taken in diversification of crops might be less feared than in middle scale farms. Additionally, large scale farms with high capital could test crop diversification and pesticide reduction on a small part of the farm to begin with, to take less risks. This system is already used by big vine producers in France, where small parts of the vineyards are converted to biodynamic vine production, this product being more and more appreciated by consumers who are willing to pay more for it. Finally, the

system might be more difficult to apply to mid-size farms as they might be committed to specialized capital equipment which would make temporal crop diversification more difficult and are too small to manage diversification in space.

A final blocking point might be the lack of interest, or maybe even the opposition, of large agrochemical companies to the proposed system. Agrochemical companies have a high influence on the agricultural sector. Today, only a few firms own a large part of the agricultural chemical market (Clapp, 2021). These firms exert an important power, more or less directly, on the way food is produced (Clapp, 2021). As they profit directly from the commercialization of chemicals, going toward a system without these chemicals as proposed in this paper could be of huge commercial challenge, but one that is consistent with competitive use of new biotechnologies with more environmentally friendly properties. As stated by Clapp (2021), these companies shape the food markets, technologies and innovation perspectives as well as policy and governance decisions. More and more, these companies are opening market branches in biological control, which actually goes in the direction of reducing the use of chemicals. However, what we propose in this paper is a way to attract and maintain natural enemies already present in the landscape (conservation biological control), with consequently no need for external inputs. It would therefore be quite optimistic to think that implementing such large-scale conservation biological programs might be enhanced by these agrochemical companies if they implement innovative environmentally compatible technologies in their market strategies.

3.2. Actions to be taken by the stakeholders to reach the next step

Diversifying crop species, and potentially crop genetics, at the landscape scale seems to have high potential for the preservation of natural




Fig. 4. Schematic representation of the agricultural network studied in relation to the economic, social and environmental pillars, including new actions to be taken by the stakeholders of the agricultural chain in the framework of landscape crop diversification. New stakeholders are highlighted in dark green and new actions are indicated with a dark grey arrow. The collaborative network for landscape design is indicated by dark green lines.

enemies and for the increase in their efficacy. Economically, the switch from a conventional system to a more diversified one will require specific attention to the markets targeted by farmers and to the possibility of providing technical support to farmers. In order to improve the success of such agricultural systems, it is important to take into consideration the point of view and the advice of different specialists.

The implementation of a system based on landscape crop diversity could be feasible if the different agricultural stakeholders are involved (Landis, 2017; Haan et al., 2021). Fig. 4 is a schematic representation of the agricultural network studied in relation to the economic, social and environmental pillars, including new actions to be taken by the stakeholders of the agricultural chain in the framework of landscape crop diversification. As mentioned earlier, new stakeholders – highlighted in dark green in Fig. 4 – need to enter in the decision making in order to optimize the proposed way to implement crop diversity schemes at a landscape scale. Economists, sociologists, geographers and advisers need to be included in a collaborative framework (Landis, 2017; Haan et al., 2021) as well as of course farmers and environmentalists already mentioned in Fig. 2. Specialized advisers could take the role of transferring the specific knowledge to farmers and these farmers should provide feedback on the proposed practices and outcomes. By implementing this new landscape design, new relations between the different agricultural stakeholders could raise. First, in order to avoid any economic losses possibly due to the changes of practices, the State could propose to the farmers a system of insurance encouraging them to take risks. Secondly, environmentalists could propose new agricultural landscape designs in collaboration with geographers, economists, sociologists and farmers in order to stick to realistic solutions. Thirdly, food retailers would inform consumers about the changes of practices in order to encourage them to change their consumption habits and help the farmers to switch from a pesticide dependent practice to a more environmentally friendly one. Within the retail network, cooperatives could be engaged in buying new crops under special market contracts in order to promote the selling of newly introduced plants in the region (Haan et al., 2021).

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgement

All the authors were supported by the project EUCLID (H2020-SFS-2014, grant number: 633999).

References

Adams, D.C., Salois, M.J., 2010. Local versus organic: a turn in consumer preferences and willingness-to-pay. *Renew. Agric. Food Syst.* 25 (4), 331–341.

Aguilera, G., Roslin, T., Miller, K., Tamburini, G., Birkhofer, K., Caballero-Lopez, B., Lindström, S.A.M., Öckinger, E., Rundlöf, M., Rusch, A., Smith, H.G., Bommarco, R., 2020. Crop diversity benefits carabid and pollinator communities in landscapes with semi-natural habitats. *J. Appl. Ecol.* 57 (11), 2170–2179.

Altieri, M.A., 1999. The ecological role of biodiversity in agroecosystems. *Agric. Ecosyst. Environ.* 74 (1–3), 19–31.

Andow, D., 1991. Vegetational diversity and arthropod population response. *Annu. Rev. Entomol.* 36 (1), 561–586.

Bailey, D., Schmidt-Entling, M.H., Eberhart, P., Herrmann, J.D., Hofer, G., Kormann, U., Herzog, F., 2010. Effects of habitat amount and isolation on biodiversity in fragmented traditional orchards. *J. Appl. Ecol.* 47 (5), 1003–1013.

Bale, J.S., van Lenteren, J.C., Bigler, F., 2008. Biological control and sustainable food production. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 363 (1492), 761–776.

Barzman, M., Lamichhane, J.R., Booi, K., Boonekamp, P., Desneux, N., Huber, L., Kudsk, P., Langrell, S.R.H., Ratnadass, A., Ricci, P., Sarah, J.L., Messean, A., 2015. Research and development priorities in the face of climate change and rapidly evolving pests. *Sustain. Agric. Rev.* 17, 1–27.

Baylis, K., Peplow, S., Rausser, G., Simon, L., 2008. Agri-environmental policies in the EU and United States: a comparison. *Ecol. Econ.* 65 (4), 753–764.

Beillouin, D., Ben-Ari, T., Makowski, D., 2019. Evidence map of crop diversification strategies at the global scale. *Environ. Res. Lett.* 14 (12), 123001.

Bernués, A., Tello-García, E., Rodríguez-Ortega, T., Ripoll-Bosch, R., Casasús, I., 2016. Agricultural practices, ecosystem services and sustainability in high nature value farmland: unraveling the perceptions of farmers and nonfarmers. *Land Use Policy* 59, 130–142.

Bianchi, F.J.J., Booij, C.J.H., Tscharntke, T., 2006. Sustainable pest regulation in agricultural landscapes: a review on landscape composition, biodiversity and natural pest control. *Proc. R. Soc. B Biol. Sci.* 273 (1595), 1715–1727.

Birner, R., Daum, T., Pray, C., 2021. Who drives the digital revolution in agriculture? A review of supply-side trends, players and challenges. *Appl. Econ. Perspect. Policy.*

Bortolotto, O.C., Júnior, M., de Oliveira, A., Hoshino, A.T., Campos, T.A., 2016. Distance from the edge of forest fragments influence the abundance of aphidophagous hoverflies (Diptera: Syrphidae) in wheat fields. *Acta Sci. Agron.* 38 (2), 157–164.

Bosem, Baillod A., Tscharntke, T., Clough, Y., Batáry, P., 2017. Landscape-scale interactions of spatial and temporal cropland heterogeneity drive biological control of cereal aphids. *J. Appl. Ecol.* 54 (6), 1804–1813.

Bouttes, M., Darnhofer, I., Martin, G., 2019. Converting to organic farming as a way to enhance adaptive capacity. *Org. Agric.* 9 (2), 235–247.

Brewer, M.J., Goodell, P.B., 2012. Approaches and incentives to implement integrated pest management that addresses regional and environmental issues. *Annu. Rev. Entomol.* 57, 41–59.

Brittain, C., Bommarco, R., Vighi, M., Settele, J., Potts, S.G., 2010. Organic farming in isolated landscapes does not benefit flower-visiting insects and pollination. *Biol. Conserv.* 143 (8), 1860–1867.

Burton, M., Rigby, D., Young, T., 1999. Analysis of the determinants of adoption of organic horticultural techniques in the UK. *J. Agric. Econ.* 50 (1), 47–63.

Chapman, J.W., Reynolds, D.R., Brooks, S.J., Smith, A.D., Woiwod, I.P., 2006. Seasonal variation in the migration strategies of the green lacewing *Chrysoperla carnea* species complex. *Ecol. Entomol.* 31 (4), 378–388.

Chifflet, R., Klein, E.K., Lavigne, C., Le Féon, V., Ricroch, A.E., Lecomte, J., Vaissiere, B.E., 2011. Spatial scale of insect-mediated pollen dispersal in oilseed rape in an open agricultural landscape. *J. Appl. Ecol.* 48 (3), 689–696.

Clapp, J., 2021. The problem with growing corporate concentration and power in the global food system. *Nat. Food* 2 (6), 404–408.

Craheix, D., Angevin, F., Doré, T., De Tourdonnet, S., 2016. Using a multicriteria assessment model to evaluate the sustainability of conservation agriculture at the cropping system level in France. *Eur. J. Agron.* 76, 75–86.

Cranfield, J.A., Magnusson, E., 2003. Canadian consumers' willingness to pay for pesticide-free food products: an ordered probit analysis. *Int. Food Agribusiness Manag. Rev.* 6 (4), 13–30.

Crawford, E., 2019. Innovative manufacturers, organic farmers team to create a win-win demand for rotational crops. In <https://www.foodnavigator-usa.com/Article/2019/02/06/Innovative-manufacturers-organic-farmers-team-to-create-a-win-win-demand-for-rotational-crops>.

Cumming, G.S., Spiesman, B.J., 2006. Regional problems need integrated solutions: pest management and conservation biology in agroecosystems. *Biol. Conserv.* 131 (4), 533–543.

Dainese, M., Schneider, G., Krauss, J., Steffan-Dewenter, I., 2017. Complementarity among natural enemies enhances pest suppression. *Sci. Rep.* 7 (1), 8172.

Degani, E., Leigh, S.G., Barber, H.M., Jones, H.E., Lukac, M., Sutton, P., Potts, S.G., 2019. Crop rotations in a climate change scenario: short-term effects of crop diversity on resilience and ecosystem service provision under drought. *Agric. Ecosyst. Environ.* 285, 106625.

Deguine, J.P., Aubertot, J.N., Flor, R.J., Lescouret, F., Wyckhuys, K.A., Ratnadass, A., 2021. Integrated pest management: good intentions, hard realities. A review. *Agron. Sustain. Dev.* 41 (3), 1–35.

Desneux, N., Decourtye, A., Delpuech, J.-M., 2007. The sublethal effects of pesticides on beneficial arthropods. *Annu. Rev. Entomol.* 52 (1), 81–106.

Eilenberg, J., Hajek, A., Lomer, C., 2001. Suggestions for unifying the terminology in biological control. *BioControl* 46 (4), 387–400.

Elkington, J., 1994. Towards the sustainable corporation: win-win-win business strategies for sustainable development. *Calif. Manag. Rev.* 36 (2), 90–100.

Evans, E.W., 2003. Searching and reproductive behaviour of female aphidophagous ladybirds 1549 (Coleoptera: Coccinellidae): a review. *Eur. J. Entomol.* 100, 1–10.

Fahrig, L., Baudry, J., Brotons, L., Burel, F.G., Crist, T.O., Fuller, R.J., Sirami, C., Siriwardena, G.M., Martin, J.L., 2011. Functional landscape heterogeneity and animal biodiversity in agricultural landscapes. *Ecol. Lett.* 14 (2), 101–112.

Fahrig, L., Girard, J., Duro, D., Pasher, J., Smith, A., Javorek, S., Tischendorf, L., 2015. Farmlands with smaller crop fields have higher within-field biodiversity. *Agric. Ecosyst. Environ.* 200, 219–234.

FAO, 2019. Can we ditch intensive farming - and still feed the human race? [online] Available at: Global Soil Partnership | Food and Agriculture Organization of the United Nations. <http://www.fao.org/global-soil-partnership/resources/highlights/detail/en/c/1179073/>

Fischbein, D., Jofré, N., Corley, J.C., 2016. A comparative analysis of host feeding and life-history traits in parasitoid wasps. *Entomol. Exp. Appl.* 159 (2), 172–180.

Fischer, R., 1980. The self image of biological farmers. An empirical sociological investigation of their motivation, way of thinking and attitude to their job [PhD thesis, ETH Zurich]. Retrieved from ETH Zurich online repository. <https://www.cabdirect.org/cabdirect/abstract/19821887484>.

Fischer, J., Lindenmayer, D.B., Manning, A.D., 2006. Biodiversity, ecosystem function, and resilience: ten guiding principles for commodity production landscapes. *Front. Ecol. Environ.* 4 (2), 80–86.

Fischer, J., Brittain, C., Klein, A.M., 2013. Biodiversity-friendly farming. *Encyclopedia of Biodiversity*, pp. 418–429.

- Forkuoh, F., Boadi, N.O., Borquaye, L.S., Afful, S., 2018. Risk of human dietary exposure to organochlorine pesticide residues in fruits from Ghana. *Sci. Rep.* 8, 16686. <https://doi.org/10.1038/s41598-018-35205-w>.
- Fornetti, A., 2019. Stakeholders' needs and expectations for an optimal implementation of IPM in agriculture. 12 septEUCLID Final Conference, Avignon. Oral Presentation.
- Gardarin, A., Plantegenest, M., Bischoff, A., Valantin-Morison, M., 2018. Understanding plant-arthropod interactions in multitrophic communities to improve conservation biological control: useful traits and metrics. *J. Pest. Sci.* 1–13.
- Gardiner, M.M., Neal, M.E.O., 2009. Landscape diversity enhances biological control of an introduced crop pest in the north-central USA. *Ecol. Appl.* 19 (1), 143–154.
- Garibaldi, L.A., Oddi, F.J., Miguez, F.E., Bartomeus, I., Orr, M.C., Jobbágy, E.G., Zhu, C.D., 2021. Working landscapes need at least 20% native habitat. *Conserv. Lett.* 14 (2), e12773.
- Gilbert, C.L., Morgan, C.W., 2010. Food price volatility. *Philos. Trans. R. Soc. B Biol. Sci.* 365 (1554), 3023–3034.
- Gillespie, M.A., Gurr, G.M., Wratten, S.D., 2016. Beyond nectar provision: the other resource requirements of parasitoid biological control agents. *Entomologia Exp. Appl.* 159 (2), 207–221.
- Global Newswire, 2019. Sustainable ag practices are a win-win. Montana organic farmers tells Congress. In <https://www.marketwatch.com/press-release/sustainable-ag-practices-are-a-win-win-montana-organic-farmer-tells-congress-2019-04-11>.
- Goldman, R.L., Thompson, B.H., Daily, G.C., 2007. Institutional incentives for managing the landscape: inducing cooperation for the production of ecosystem services. *Ecol. Econ.* 64 (2), 333–343.
- Guichard, L., Dedieu, F., Jeuffroy, M.H., Meynard, J.M., Reau, R., Savini, I., 2017. Ecophyto, the French action plan to reduce pesticide use: a failure analyses and reasons for hoping. *Cah. Agric.* 26 (1).
- Gurr, G.M., Wratten, S.D., Landis, D.A., You, M., 2017. Habitat management to suppress pest populations: progress and prospects. *Annu. Rev. Entomol.* 62, 91–109.
- Haan, N.L., Zhang, Y., Landis, D.A., 2020. Predicting landscape configuration effects on agricultural pest suppression. *Trends Ecol. Evol.* 35 (2), 175–186.
- Haan, N.L., Iuliano, B.G., Gratton, C., Landis, D.A., 2021. Designing agricultural landscapes for arthropod-based ecosystem services in North America. *The Future of Agricultural Landscapes*, Part II. 191.
- Haile, M.G., Wossen, T., Tesfaye, K., von Braun, J., 2017. Impact of climate change, weather extremes, and price risk on global food supply. *Econ. Disasters Clim. Chang.* 1 (1), 55–75.
- InHassel, T., 2014. Small vs. large: which size farm is better for the planet? https://www.washingtonpost.com/lifestyle/food/small-vs-large-which-size-farm-is-better-for-the-planet/2014/08/29/ac2a3dc8-2e2d-11e4-994d-202962a9150c_story.html?noredirect=on&utm_term=.5b693ce0d0f6
- Hatt, S., Francis, F., Xu, Q., Wang, S., Osawa, N., 2020. Perennial flowering strips for conservation biological control of insect pests: from picking and mixing flowers to tailored functional diversity. *Integrative Biological Control*, pp. 57–71 https://doi.org/10.1007/978-3-030-44838-7_4.
- He, H.M., Liu, L.N., Munir, S., Bashir, N.H., Yi, W.A.N.G., Jing, Y.A.N.G., Li, C.Y., 2019. Crop diversity and pest management in sustainable agriculture. *J. Integr. Agric.* 18 (9), 1945–1952.
- Holzschuh, A., Steffan-Dewenter, I., Tscharntke, T., 2010. How do landscape composition and configuration, organic farming and fallow strips affect the diversity of bees, wasps and their parasitoids? *J. Anim. Ecol.* 79 (2), 491–500.
- Hulme, P.E., 2003. Biological invasions: winning the science battles but losing the conservation war? *Oryx* 37 (2), 178–193.
- Ibrahim, I.H.M., Gilfoyle, L., Reynolds, R., Voulvoulis, N., 2019. Integrated catchment management for reducing pesticide levels in water: engaging with stakeholders in East Anglia to tackle metaldehyde. *Sci. Total Environ.* 656, 1436–1447.
- Iuliano, B., Gratton, C., 2020. Temporal resource (dis) continuity for conservation biological control: from field to landscape scales. *Front. Sustain. Food Syst.* 4, 127.
- Jauker, F., Diekoetter, T., Schwarzbach, F., Wolters, V., 2009. Pollinator dispersal in an agricultural matrix: opposing responses of wild bees and hoverflies to landscape structure and distance from main habitat. *Landsc. Ecol.* 24 (4), 547–555.
- Jeaneret, P., Lüscher, G., Herzog, F., 2012. Methods for assessing biodiversity indicators at farm scale. *Biodiversity Indicators for European Farming Systems*. 25.
- Jokanovic, M., 2018. Neurotoxic effects of organophosphorus pesticides and possible association with neurodegenerative diseases in man: a review. *Toxicology* 410, 125–131.
- Jonsson, M., Wratten, S.D., Landis, D.A., Gurr, G.M., 2008. Recent advances in conservation biological control of arthropods by arthropods. *Biol. Control* 45 (2), 172–175.
- Karp, D.S., Chaplin-Kramer, R., Meehan, T.D., Martin, E.A., DeClerck, F., Grab, H., Gratton, C., Hunt, L., Larsen, A.E., Martínez-Salinas, A., O'Rourke, M.E., 2018. Crop pests and predators exhibit inconsistent responses to surrounding landscape composition. *Proc. Natl. Acad. Sci.* 115 (33), E7863–E7870.
- Kheirodin, A., Cárcamo, H.A., Costamagna, A.C., 2020. Contrasting effects of host crops and crop diversity on the abundance and parasitism of a specialist herbivore in agricultural landscapes. *Landsc. Ecol.* 35 (5), 1073–1087.
- Kilpatrick, S., Rosenblatt, T., 1998. Information vs training: issues in farmer learning. *J. Agric. Educ. Ext.* 5 (1), 39–51.
- Kip-Tot, E., Lukuyu, B., Franzel, S., Place, F., 2011. Farmers teaching farmers: challenges and opportunities of using volunteer farmers in technology dissemination. *International Conference on Innovations in Extension and Advisory Services*, November, pp. 15–18.
- Koricheva, J., Hayes, D., 2018. The relative importance of plant intraspecific diversity in structuring arthropod communities: a meta-analysis. *Funct. Ecol.* 32 (7), 1704–1717.
- Lacey, L.A., Frutos, R., Kaya, H.K., Vail, P., 2001. Insect pathogens as biological control agents: do they have a future? *Biol. Control* 21 (3), 230–248.
- Landis, D.A., 2017. Designing agricultural landscapes for biodiversity-based ecosystem services. *Basic Appl. Ecol.* 18, 1–12.
- Larsen, A.E., Noack, F., 2021. Impact of local and landscape complexity on the stability of field-level pest control. *Nat. Sustain.* 4 (2), 120–128.
- Leach, A.W., Mumford, J.D., 2008. Pesticide environmental accounting: a method for assessing the external costs of individual pesticide applications. *Environ. Pollut.* 151, 139–147.
- Letourneau, D.K., Jedlicka, J.A., Bothwell, S.G., Moreno, C.R., 2009. Effects of natural enemy biodiversity on the suppression of arthropod herbivores in terrestrial ecosystems. *Annu. Rev. Ecol. Evol. Syst.* 40, 573–592.
- Letourneau, D.K., Armbrrecht, I., Rivera, B.S., Lerma, J.M., Carmona, E.J., Daza, M.C., Escobar, S., Galindo, V., Gutiérrez, C., López, S.D., Mejía, J.L., 2011. Does plant diversity benefit agroecosystems? A synthetic review. *Ecol. Appl.* 21 (1), 9–21.
- Lew, S., Lew, M., Szarek, J., Mieszczynski, T., 2009. Effect of pesticides on soil and aquatic environmental microorganisms - a short review. *Fresenius Environ. Bull.* 18 (8), 1390–1395.
- Li, C., Yang, J., He, X., Zhu, S., Zhu, Y., 2020. Agricultural Biodiversity for Crop Disease and Pest Management. *Integrative Biological Control*. Springer, Cham, pp. 117–132.
- Lin, B.B., 2011. Resilience in agriculture through crop diversification: adaptive management for environmental change. *Bioscience* 61 (3), 183–193.
- Liu, B., Yang, L., Yang, Y., Lu, Y., 2016. Influence of landscape diversity and composition on the parasitism of cotton bollworm eggs in maize. *PLoS One* 11 (2).
- Liu, B., Yang, L., Zeng, Y., Yang, F., Yang, Y., Lu, Y., 2018. Secondary crops and non-crop habitats within landscapes enhance the abundance and diversity of generalist predators. *Agric. Ecosyst. Environ.* 258, 30–39.
- Liu, T., Wang, X., Chen, D., Li, Y., Wang, F., 2018. Growth, reproduction and biochemical toxicity of chlorantraniliprole in soil on earthworms (*Eisenia fetida*). *Ecotoxicol. Environ. Saf.* 150, 18–25.
- Lundgren, J.G., Hesler, L.S., Clay, S.A., Fausti, S.F., 2013. Insect communities in soybeans of eastern South Dakota: the effects of vegetation management and pesticides on soybean aphids, bean leaf beetles, and their natural enemies. *Crop Prot.* 43, 104–118.
- Martin, E.A., Seo, B., Park, C.R., Reineking, B., Steffan-Dewenter, I., 2016. Scale-dependent effects of landscape composition and configuration on natural enemy diversity, crop herbivory, and yields. *Ecol. Appl.* 26 (2), 448–462.
- Martin, E.A., Dainese, M., Clough, Y., Baldi, A., Bommarco, R., Gagic, V., Steffan-Dewenter, I., 2019. The interplay of landscape composition and configuration: new pathways to manage functional biodiversity and agroecosystem services across Europe. *Ecol. Lett.* 22 (7), 1083–1094.
- Matthews, A., 2013. Greening agricultural payments in the EU's common agricultural policy. *Bio-based Appl. Econ.* 2 (1), 1–27.
- Meynard, J.M., Messéan, A., Charlier, A., Charrier, F., Le Bail, M., Magrini, M.B., Savini, I., 2013. Freins et leviers à la diversification des cultures: étude au niveau des exploitations agricoles et des filières. *Ocl* 20 (4), D403.
- Meynard, J.M., Charrier, F., Le Bail, M., Magrini, M.B., Charlier, A., Messéan, A., 2018. Socio-technical lock-in hinders crop diversification in France. *Agron. Sustain. Dev.* 38 (5), 1–13.
- Monticelli, L.S., Nguyen, L.T., Amiens-Desneux, E., Luo, C., Lavoie, A.V., Gatti, J.L., Desneux, N., 2019. The preference-performance relationship as a means of classifying parasitoids according to their specialization degree. *Evol. Appl.* 12, 1626–1640.
- Morel, K., Revoyron, E., San Cristobal, M., Baret, P.V., 2020. Innovating within or outside dominant food systems? Different challenges for contrasting crop diversification strategies in Europe. *PLoS One* 15 (3), e0229910.
- Nagayets, O., 2005. Small farms: current status and key trends. *The Future of Small Farms*. 355.
- Nelson, K.S., Burchfield, E.K., 2021. Landscape complexity and US crop production. *Nat. Food* 2 (5), 330–338.
- Nicholls, C.I., Altieri, M.A., 2013. Plant biodiversity enhances bees and other insect pollinators in agroecosystems. A review. *Agron. Sustain. Dev.* 33 (2), 257–274.
- Nicholson, C.C., Williams, N.M., 2021. Cropland heterogeneity drives frequency and intensity of pesticide use. *Environ. Res. Lett.* 16 (7), 074008.
- Olsson, R., 1988. Management for success in modern agriculture. *Eur. Rev. Agric. Econ.* 15 (2–3), 239–259.
- Osawa, N., 2000. Population field studies on the aphidophagous ladybird beetle *Harmonia axyridis* (Coleoptera: Coccinellidae): resource tracking and population characteristics. *Popul. Ecol.* 42 (2), 115–127.
- Otway, S.J., Hector, A., Lawton, J.H., 2005. Resource dilution effects on specialist insect herbivores in a grassland biodiversity experiment. *J. Anim. Ecol.* 74 (2), 234–240.
- Peattie, K., 2001. Towards sustainability: the third age of green marketing. *Mark. Rev.* 2 (2), 129–146.
- Peña, J.E., 2013. *Potential Invasive Pests of Agricultural Crops*. vol. 3. CABI.
- Perovic, D., Gámez-Virués, S., Börschig, C., Klein, A.M., Krauss, J., Rothenwöhler, C., Erasmí, S., Tschamtké, T., Westphal, C., 2015. Configurational landscape heterogeneity shapes functional community composition of grassland butterflies. *J. Appl. Ecol.* 52 (2), 505–513.
- Pingali, P.L., 2012. Green revolution: impacts, limits, and the path ahead. *Proc. Natl. Acad. Sci.* 109 (31), 12302–12308.
- Potter, C., Lobley, M., 1993. Helping small farms and keeping Europe beautiful: a critical review of the environmental case for supporting the small family farm. *Land Use Policy* 10 (4), 267–279.
- Provost, D., Gruber, A., Pierre, L.E., Jaffré, A., Loyant, V., Loiseau, H., Vital, A., Brochard, P., Baldi, I., 2007. Brain tumors and exposure to pesticides: a case-control study in southwestern France. *Occup. Environ. Med.* 64 (8), 509–514.
- Rand, T.A., Tschamtké, T., 2007. Contrasting effects of natural habitat loss on generalist and specialist aphid natural enemies. *Oikos* 116 (8), 1353–1362.
- Raynaud, E., Sauvée, L., Valceschini, E., 2009. Aligning branding strategies and governance of vertical transactions in agri-food chains. *Ind. Corp. Chang.* 18 (5), 835–868.
- Redlich, S., Martin, E.A., Steffan-Dewenter, I., 2018. Landscape-level crop diversity benefits biological pest control. *J. Appl. Ecol.* <https://doi.org/10.1111/1365-2664.13126>.

- Rigby, D., Young, T., Burton, M., 2001. The development of and prospects for organic farming in the UK. *Food Policy* 26 (6), 599–613.
- InRoh, C., 2013. How do spiders move? Cornell University's naturalist outreach. <https://cpb-us-e1.wpmucdn.com/blogs.cornell.edu/dist/7/3643/files/2013/09/How-Do-Spiders-Move-1bpbzvbv.pdf>.
- Root, R.B., 1973. Organization of a plant-arthropod association in simple and diverse habitats: the fauna of collards (*Brassica oleracea*). *Ecol. Monogr.* 43 (1), 95–124.
- Rusch, A., Bommarco, R., Jonsson, M., Smith, H.G., Ekbom, B., 2013. Flow and stability of natural pest control services depend on complexity and crop rotation at the landscape scale. *J. Appl. Ecol.* 50 (2), 345–354.
- Rusch, A., Chaplin-Kramer, R., Gardiner, M.M., Hawro, V., Holland, J., Landis, D., Bommarco, R., 2016. Agricultural landscape simplification reduces natural pest control: a quantitative synthesis. *Agric. Ecosyst. Environ.* 221, 198–204.
- Schaller, N., 2012. La diversification des assolements en France: intérêts, freins et enjeux. Analyse. Centre d'études et de prospective 4pp.
- Schellhorn, N.A., Bianchi, F.J.J.A., Hsu, C.L., 2014. Movement of entomophagous arthropods in agricultural landscapes: links to pest suppression. *Annu. Rev. Entomol.* 59, 559–581.
- Schellhorn, N.A., Gagic, V., Bommarco, R., 2015. Time will tell: resource continuity bolsters ecosystem services. *Trends Ecol. Evol.* 30 (9), 524–530.
- Sharpley, R., Vass, A., 2006. Tourism, farming and diversification: an attitudinal study. *Tour. Manag.* 27 (5), 1040–1052.
- Shreck, A., Getz, C., Feenstra, G., 2006. Social sustainability, farm labor, and organic agriculture: findings from an exploratory analysis. *Agric. Hum. Values* 23 (4), 439–449.
- Sirami, C., Gross, N., Bailod, A.B., Bertrand, C., Carrié, R., Hass, A., Fahrig, L., 2019. Increasing crop heterogeneity enhances multitrophic diversity across agricultural regions. *Proc. Natl. Acad. Sci.* 116 (33), 16442–16447.
- Sirieix, L., Delanchy, M., Remaud, H., Zepeda, L., Gurviez, P., 2013. Consumers' perceptions of individual and combined sustainable food labels: a UK pilot investigation. *Int. J. Consum. Stud.* 37 (2), 143–151.
- Slotterback, C.S., Runck, B., Pitt, D.G., Kne, L., Jordan, N.R., Mulla, D.J., Zenger, C., Reichenbach, M., 2016. Collaborative geodesign to advance multifunctional landscapes. *Landsc. Urban Plan.* 156, 71–80.
- Smith-Spangler, C., Brandeau, M.L., Hunter, G.E., Bavinger, J.C., Pearson, M., Eschbach, P.J., et al., 2012. Are organic foods safer or healthier than conventional alternatives?: A systematic review. *Ann. Intern. Med.* 157 (5), 348–366.
- Snyder, L.D., Gómez, M.I., Mudrak, E.L., Power, A.G., 2021. Landscape-dependent effects of varietal mixtures on insect pest control and implications for farmer profits. *Ecol. Appl.* 31 (2), e02246.
- Stallman, H.R., James Jr., H.S., 2015. Determinants affecting farmers' willingness to cooperate to control pests. *Ecol. Econ.* 117, 182–192.
- Stiling, P., Cornelissen, T., 2005. What makes a successful biocontrol agent? A meta-analysis of biological control agent performance. *Biol. Control* 34 (3), 236–246.
- Symondson, W.O.C., Sunderland, K.D., Greenstone, M.H., 2002. Can generalist predators be effective biocontrol agents? *Annu. Rev. Entomol.* 47 (1), 561–594.
- Taning, C.N., Arpaia, S., Christiaens, O., Dietz-Pfeilstetter, A., Jones, H., Mezzetti, B., Smagghe, G., 2020. RNA-based biocontrol compounds: current status and perspectives to reach the market. *Pest Manag. Sci.* 76 (3), 841–845.
- Tscharntke, T., Rand, T.A., Bianchi, F.J., 2005. The landscape context of trophic interactions: insect spillover across the crop–non crop interface. *Annales Zoologici Fennici*. Finnish Zoological and Botanical Publishing Board, pp. 421–432.
- Tscharntke, T., Tylianakis, J.M., Wade, M.R., Wratten, S.D., Bengtsson, J., Kleijn, D.A.V.I.D., 2007. Insect conservation in agricultural landscapes. *Insect Conserv. Biol.* 16, 383–404.
- Tuda, M., Shima, K., 2002. Relative importance of weather and density dependence on the dispersal and on-plant activity of the predator *Orius minutus*. *Popul. Ecol.* 44 (3), 0251–0257.
- Valbuena, D., Verburg, P.H., Bregt, A.K., Ligtenberg, A., 2010. An agent-based approach to model land-use change at a regional scale. *Landsc. Ecol.* 25 (2), 185–199.
- Van der Valk, H., Koomen, I., Blacquiere, T., Van der Steen, J.J.M., Roessink, I., 2011. Aspects Determining the Risk of Pesticides to Wild Bees: Risk Profiles for Focal Crops on Three Continents. Food and Agriculture Organization of the United Nations.
- Van Rijn, P.C., Kooijman, J., Wäckers, F.L., 2013. The contribution of floral resources and honeydew to the performance of predatory hoverflies (Diptera: Syrphidae). *Biol. Control* 67 (1), 32–38.
- Vasseur, C., Joannon, A., Aviron, S., Burel, F., Meynard, J.M., Baudry, J., 2013. The cropping systems mosaic: how does the hidden heterogeneity of agricultural landscapes drive arthropod populations? *Agric. Ecosyst. Environ.* 166, 3–14.
- Van Zanten, B.T., Verburg, P.H., Espinosa, M., Gomez-y-Paloma, S., Galimberti, G., Kantelhardt, J., ... Viaggi, D., 2014. European agricultural landscapes, common agricultural policy and ecosystem services: a review. *Agron. Sustain. Dev.* 34 (2), 309–325.
- Velki, M., Weltmeyer, A., Seiler, T.B., Hollert, H., 2019. Acute toxicities and effects on multixenobiotic resistance activity of eight pesticides to the earthworm *Eisenia andrei*. *Environ. Sci. Pollut. Res.* 26 (5), 4821–4832.
- Villeneuve-Chasset, J., 2006. Etude de la Bio-écologie des Névroptères dans une perspective de lutte biologique par conservation [PhD thesis, Angers University]. Retrieved from Angers University online repository. <https://tel.archives-ouvertes.fr/tel-00198786>.
- Wagner, D.L., 2020. Insect declines in the Anthropocene. *Annu. Rev. Entomol.* 65, 457–480.
- Wahlang, B., 2018. Exposure to persistent organic pollutants: impact on women's health. *Rev. Environ. Health* 33 (4), 331–348.
- Wan, N.F., Zheng, X.R., Fu, L.W., Kiær, L.P., Zhang, Z., Chaplin-Kramer, R., Li, B., 2020. Global synthesis of effects of plant species diversity on trophic groups and interactions. *Nat. Plants* 6 (5), 503–510.
- Wang, X.G., Keller, M.A., 2003. Patch time allocation by the parasitoid *Diadegma semiclausum* (Hymenoptera: Ichneumonidae). I. Effect of interpatch distance. *J. Insect Behav.* 16 (2), 279–293.
- Wezel, A., Goette, J., Lagneaux, E., Passuello, G., Reisman, E., Rodier, C., Turpin, G., 2018. Agroecology in Europe: research, education, collective action networks, and alternative food systems. *Sustainability* 10 (4), 1214.
- Wilson, C., Tisdell, C., 2001. Why farmers continue to use pesticides despite environmental, health and sustainability costs. *Ecol. Econ.* 39 (3), 449–462.
- Woodcock, B.A., Bullock, J.M., Shore, R.F., Heard, M.S., Pereira, M.G., Redhead, J., et al., 2017. Country-specific effects of neonicotinoid pesticides on honey bees and wild bees. *Science* 356 (6345), 1393–1395.
- Wratten, S.D., Gillespie, M., Decourtye, A., Mader, E., Desneux, N., 2012. Pollinator habitat enhancement: benefits to other ecosystem services. *Agric. Ecosyst. Environ.* 159, 112–122.
- Zhao, H., Li, J., Guo, L., Wang, K., 2021. Crop diversity at the landscape level affects the composition and structure of the vegetation-dwelling arthropod communities in naked oat (*Avena chinensis*) fields. *Int. J. Environ. Res. Public Health* 18 (1), 30.
- Ziska, L.H., Blumenthal, D.M., Runion, G.B., Hunt, E.R., Diaz-Soltero, H., 2011. Invasive species and climate change: an agronomic perspective. *Clim. Chang.* 105 (1), 13–42.